



CLAIMS

1. Process for preparing block polymers of general formula (IA) or (IB):



5

in which formulae:

- R' represents:

- an optionally substituted alkyl, acyl, aryl, alkene or alkyne group (i), or
 - 10 • an optionally substituted or aromatic, saturated or unsaturated, carbocycle (ii), or
 - an optionally substituted or aromatic, saturated or unsaturated, heterocycle (iii),
- it being possible for these groups and rings (i),
- 15 (ii) and (iii) to be substituted with substituted phenyl groups, substituted aromatic groups, or groups: alkoxycarbonyl or aryloxycarbonyl (-COOR), carboxyl (-COOH), acyloxy (-O₂CR), carbamoyl (-CONR₂), cyano (-CN), alkylcarbonyl, alkylarylcarbonyl,

5

10

15

~~-R² and R³, which are identical or different, represent:~~

- 20

it being possible for these groups and rings (i),

25

- . substituted phenyl groups or substituted aromatic groups,
. groups: alkoxycarbonyl or aryloxy carbonyl



(-COOR), carboxyl (-COOH), acyloxy (-O₂CR),
 carbamoyl (-CONR₂), cyano (-CN), alkylcarbonyl,
 alkylarylcabonyl, arylcarbonyl,
 arylalkylcarbonyl, phthalimido, maleimido,
 5 succinimido, amidino, guanidimo, hydroxyl (-OH),
 amino (-NR₂), halogen, allyl, epoxy, alkoxy (-OR),
 S-alkyl, S-aryl,
 . groups having a hydrophilic or ionic character,
 such as the alkali metal salts of carboxylic
 10 acids, the alkali metal salts of sulphonic acid,
 polyalkylene oxide chains (PEO, PPO), cationic
 substituents (quaternary ammonium salts),
 R representing an alkyl or aryl group,
 and, for at least R² or R³, these groups and rings (i),
 15 (ii) and (iii) induce a delocalizing or electron-
 withdrawing effect with respect to the electron density
 of the nitrogen atom to which R² and R³ are linked,
 - V, V', W and W', which are identical or different,
 represent: H, an alkyl group or a halogen,
 20 - X, X', Y and Y', which are identical or different,
 represent H, a halogen or a group chosen from R', OR',
 OCOR', NHCOH, OH, NH₂, NHR', N(R')₂, (R')₂N⁺O⁻, NHCOR',
 CO₂H, CO₂R', CN, CONH₂, CONHR' or CONR'₂, in which R' is
 25 chosen from alkyl, aryl, aralkyl, alkaryl, alkene or
 organosilyl groups, optionally perfluorinated and
 optionally substituted with one or more carboxyl,
 epoxy, hydroxyl, alkoxy, amino, halogen or sulphonic
 groups,

001160:000000

- **a** and **b**, which are identical or different, are equal to 0 or 1,
- **m** and **n**, which are identical or different, are greater than or equal to 1 and, when one or other is greater than 1, the individual repeat units are identical or different,

in which process the following are brought into contact with each other:

- an ethylenically unsaturated monomer of formula:
 $CYY' (=CW-CW')=CH_2$,
- a precursor compound of general formula (IIA) or (IIB):



- in which Z, X, X', V, V', R¹, R² and R³ have the same meaning, and b and n the same value, as previously;
- a radical polymerization initiator.

2. Process according to claim 1, characterized in that the ethylenically unsaturated monomer is chosen from: styrene or its derivatives, butadiene,

chloroprene, (meth)acrylic esters and vinyl nitriles.

3. Process according to claim 1 or 2, characterized in that, in compounds of formula (IA) and (IIA), the ring Z is a ring based on carbon atoms.

5 4. Process according to any one of the preceding claims, characterized in that the ring Z includes at least one heteroatom other than the nitrogen which links the ring Z to $-C(=S)S$, this heteroatom being chosen from O, S, N and/or P.

10 5. Process according to any one of the preceding claims, characterized in that, in the compounds of formula (IA) and (IIA), the ring Z is an aromatic ring.

6. Process according to any one of the preceding claims, characterized in that, in the compounds of
15 formula (IA) and (IIA), the ring Z comprises at least one of the following functional groups: carbonyl ($C=O$), SO_2 , POR'' , R'' representing an alkyl, aryl, OR, SR or NR_2 group, where the R is identical or different and represents an alkyl or aryl group.

20 7. Process according to any one of the preceding claims, characterized in that, in the compounds of formula (IA) and (IIA), the ring Z is substituted with at least one of the following groups: alkyl, aryl, alkoxycarbonyl or aryloxycarbonyl ($-COOR$), carboxyl
25 ($-COOH$), acyloxy ($-O_2CR$), carbamoyl ($-CONR_2$), cyano ($-CN$), alkylcarbonyl, alkylarylcarbonyl, arylcarbonyl, arylalkylcarbonyl, phthalimido, maleimido, succinimido, amidino, guanidimo, hydroxyl ($-OH$), amino ($-NR_2$),




001160-0523560

halogen, allyl, epoxy, alkoxy (-OR), S-alkyl, S-aryl, groups having a hydrophilic or ionic character, such as the alkali metal salts of carboxylic acids or the alkali metal salts of sulphonic acid, polyalkylene
 5 oxide chains (PEO, PPO), cationic substituents (quaternary ammonium salts), R representing an alkyl or aryl group.



8. Process according to any one of the preceding claims, characterized in that, in the compounds of
 10 formula (IA) and (IIA), the ring Z is substituted with at least one carbocycle or a heterocycle, this being optionally aromatic and/or substituted.

9. Process according to the preceding claim, characterized in that, in the compounds of formula (IA)
 15 and (IIA), the ring Z and its cyclic substituent have two common atoms.

10. Process according to any one of the preceding claims, characterized in that the ring Z is chosen from one of the following rings:




001150 052350




- 10 11. Process according to claim 1, characterized
in that, in the case of the compounds of formula (IB)
and (IIB), R^2 and/or R^3 exert a π withdrawing effect.
12. Process according to claim 11, characterized
in that R^2 and/or R^3 represent a carbonyl or
15 (hetero)aromatic group.
13. Process according to claim 1, characterized
in that, in the case of the compounds of formula (IB)
and (IIB), R^2 and/or R^3 exert a Σ withdrawing effect.
14. Process according to claim 1 or 13,
20 characterized in that R^2 and/or R^3 represent an alkyl
group substituted with electron-withdrawing groups.
15. Process according to any one of the preceding
claims, characterized in that R^1 is chosen from the
groups:


81




5




10



15




20



25


16. Process according to any one of claims 1 to 10 and 15 to 6, characterized in that the precursor


compound of general formula (IIA) is a polymer and in that the said polymer comes from the radical polymerization of an ethylenically unsaturated monomer of formula: $CXX' (=CV-CV')_n=CH_2$, during which the said monomer is brought into contact with a radical polymerization initiator and a compound of general formula (IIIA) or (IVA):




p. being between 2 and 10.

17. Process according to the preceding claim, characterized in that the compound of formula (IIIA) is chosen from compounds of the following formulae:







5



10






15




20



25

001160-000000





18. Process according to any one of claims 1 and 11 to 16, characterized in that the precursor compound of general formula (IIB) is a polymer and in that the said polymer comes from the radical polymerization of an ethylenically unsaturated monomer of formula: $CXX' (=CV-CV')_n=CH_2$, during which the said monomer is brought into contact with a radical polymerization initiator and a compound of general formula (IIIB), (IVB) or (VB):






p being between 2 and 10.


- 10 19. Process according to the preceding claim, characterized in that the compound of formula (IIIB) is chosen from the compounds of the following formulae:



001160-0628500



(K)



(L)

20. Process for preparing multiblock polymers, characterized in that the implementation of the process according to one of claims 1 to 15 is repeated at least once, using:

- different monomers from those of the previous implementation, and

- instead of the precursor compound of formula (IIA) or (IIB), the block polymer coming from the previous implementation.

21. Block polymer capable of being obtained by the process according to one of the preceding claims.

22. Block polymer according to the preceding claim, characterized in that it has a polydispersity index of at most 2.

23. Block polymer according to the preceding claim, characterized in that it has a polydispersity index of at most 1.5.

00760-0623560

- polystyrene/polymethyl acrylate,
- polystyrene/polyethyl acrylate,
- polystyrene/poly(*tert*-butyl acrylate),
- polyethyl acrylate/polyvinyl acetate,
- polybutyl acrylate/polyvinyl acetate,
- poly(*tert*-butyl acrylate)/polyvinyl acetate.

26. Polymer according to claim 25, characterized in that it has a polydispersity index of at most 2.

28. Compound of general formula (IIA) or (IIB), characterized in that it has a polydispersity index of at most 2.

Mod A2

Sold B6

Add 27